BIOLOGY (231)

4.4.1 Biology Paper 1 (231/1)

1.(a)	Is when the rate of water loss is more than the rate of absorption and the plant droop	ps;	
(L)			
(b)	The rate of active transport increases with increase intemperature up to the optimun temperature;	II	1 mark
	Further increase in temperature slows down the rate of active transport until it stops denatures enzymes;	because it	1 mark
2.(a)	Animal cell;		1 mark
(b)	- Has cell membrane only/has no cell wall;		
	- Has numerous small vacuoles;		
	- Has central nucleus;	Max.	2 marks
(c)	Consists of many similar cells performing the same function;		1 mark
3.(a)	Have mammary glands; have external ears/pinna;		
	Body covered with fur/hair;	Max.	2 marks
(b)	Class;		1 mark
4.(a)	Lubrication; Protection;		2 marks
(b)	Young people are more active; requiring more energy;/		
	Older people are less active; requiring less energy;		2 marks
5.	As the cell gains water by osmosis; the sap/cellvacuole enlarges; pushing the cytople	lasm	
	outwards; exerting pressure on the cell wall;	Any 3	3 marks
6.	6000(μm)		
	55 (cells) ; 109μm;		2 marks
7.(a)	Water molecules cling to each other maintaining a continuous column of water/prev	venting the	
	break of water column;		1 mark
(b)	Water molecules cling to the sides of the xylem vessel walls;		1 mark
8.	1(a) - Leaf with serrated margin go to 2;		
	(b) - Leaf with smooth margin go to;		2 marks
9.	Presence of myelin sheath for insulation/increases transmission; Axon impulses;	for transmissio	on of
	Large cell body controls activites of cell; Nerve endings/dendrites receives impulse	s from	


	Node of Ranvier speeds up impulse transmission.			
(b)	Inner membrane highly folded/cristae to increase S A for attachment of (respiratory) e	enzym	es.	4 marks
10.	Cells loosely arranged; to facilitate air circulation;			
	Cells have moist surfaces; to dissolve respiratory gases;			
				2 marks
11.	Can receive blood from any donor/ universal recepient;			1 mark
12. (a)	(i) Arachnida;			1 mark
12. (a)	(ii) Spider/scorpion/tick/mite;			1 mark
(b)	Protoctista/protista;			1 mark
13.	Autotrophic nutrition; show alternation of generation;			
	Limited movement;			
	Limited excretory products/unspecialized respiratory structures;			2 1
	Localised growth;			2 marks
14.	Alcohol/ethanol; Carbon (IV) oxide; Energy/Adenosine Triphosphate;			3 marks
15.	To increase supply of oxygen to the tissues;The oxygen is used to oxidize lactic acid (to carbon (IV) oxide, water and energy);			2 marks
16.	Protogyny; protandry; Dioecious; Dichogamy;			2 marks
10.	Self sterility/incompatibility; Heterostyly;			
	Presence of structures/substances to attract agents of pollination;		Max.	3 marks
	,			
17.	Ovary /Anther;			1 mark
18.	- Acrosome/Lysosome contain enzyme to digest membrane of the ovum;			
	- Numerous mitochondria to provide energy for movement;			
	- Long tail for faster movement;		Max.	2 marks
19.	- Embryo not fully developed;			
	- Chemical inhibitors/presence of abscisic acid;			
	- Hard/impermeable testa/seed coat;			2 1
	- Low hormones/low enzymes concentration;	Max.		3 marks
20.	Genetically acquired beneficial characteristics which occur spontaneously; are			
	perpetuated through reproduction;			2 marks
21.(a)	Continents existed as one large Landmass/Pangea/Laurasian and Gondwana Land; Present continents drifted from it leading to isolation of organisms; organisms in each continent	h		
	evolved along different lines hence emergence of new species;			3 marks

receptors cells;


(b)	Emergence of new life/species/organisms from pre-existing simple forms, gradually over a long period of time, to present complex forms;	1 mark
22.(a)	Thigmotropism/Haptotropism;	1 mark
(b)	Part of the tendril in contact with support causes migration of auxins to the opposite side; leading to faster cell division/growth on the side not in contact with the support; This causes the tendril to curl around the support;	
		3 marks
23.	Use of biconcave/concave lens/divergent lens; to diverge the rays and make image be focussed on the retina;	2 marks
24.	 Contains antibodies that defend the body from foreign antigens; Has white blood cells that produce antibodies/while blood cells engulf antigens; Has platelets that initiate blood clotting to prevent excessive bleeding at an open wound/ prevent entry of pathogens; 	
25.	- Thin and long to allow for capillarity;	3 marks
23.	- Walls lignified to strengthen the stem/to prevent collapse of vessels;	
	- Have bordered pits to allow for exchange of materials;	2 1
	Max.	2 marks
26.(a)	Genes inherited along with the sex chromosomes;	1 mark
(b)	Haemophilia; hairy ears/pinna/nose; colour blindness/red green; blue-green colour blindness; Muscular diastrophy; baldness	
27.(a)	Complete metamorphosis - eggs hatch into larvae while in incomplete metamorphosis hatch into nymphs which resemble the adult;	2 marks
	Complete metamorphosis has four stages; egg, larvae, pupa and adult while an	
	incomplete metamorphosis has three stages; egg, nymph and adult.	2 marks
(b)	To allow for growth of the insect;	1 mark
28. (a)	Ligaments; synovial fluid; synovial membrane; articular cartilage; synovial capsule; a bone with rounded head fitting into a cavity of another bone;	2 1
(b)	Max. (i) Atlas; (ii) Axis allows movement in all planes;	2 marks 2 marks
29.	- Form joints with the legs to make walking possible;	1 mark
	Provide large surface area for attachment of muscles;Offers support (to the body weight)	1 mark
30.	Absorption of water; support; Opening and aloning of stomate:	
	Opening and closing of stomata; Feeding in insectivorous/plants;	2 marks

4.4.2 Biology Paper 2 (231/2)

1.	(a)	(i)	В	Seta/stalk;		1 mark
			d	Rhizoid;		1 mark
		(ii)	A	Production of spores/sporulation;		1 mark
			C	Photosynthesis;		1 mark
	(b)	(i)	Arthr	opoda;		1 mark
		(ii)	-	Segmented body;		
			-	Jointed appendages;		
			-	Presence of exoskeleton		3 marks
2.	(a)	E	Semi	circular canals;		
		F	Oval	window/Fenestra ovalis/Fenestra vestibuli;		
		G	Coch	lea;		3 marks
	(b)	(i)	Lined bodie	I with hair/secretion of wax/(has glands that secrete wax) to es;	trap foreig	gn
				ow/tubular/tube; to direct sound waves to the ear drum/tympa brane;	anum/tymp	oanic
			mem	orane,	(max)	(2 marks
		(ii)	Small	I/form a lever system/solid; to amplify (sound) vibrations;		(2 marks)
	(c)	Deafr	ness/abs	ence of pinna/ vertigo/tinnitus;	(max)	(1 mark
3.	(a)	(i)	Provi photo	des energy needed to split water molecules into oxygen and	hydrogen/	1
			-	des energy for formation of ATP molecules (which is used i	n dark stac	re)
			11011	des energy for formation of 1111 molecules (which is used i	ii dark stag	(1 mark)
		(ii)	Comb	pines with hydrogen ions to make glucose;		(1 mark)
		(iii)	Used	to trap light energy;		(1 mark)
	(b)	(i)	Starc	h;		
		(ii)	Prote	in;		(2 marks)
	(c)	(i)		of vitamin B1/thiamine;		(1 mark)
		(ii)	- Stu	nted growth;		
				alysis of legs/arms/limbs/damage to peripheral nerves;		
			- Hea	art failure		
			- Sw	elling of feet/oedema		
			- Gas	strointestinal disturbances/loss of appetite/sonstipation/diarr	hoea/vomi	ting;
				ight loss/muscle wasting		
			- Pal	e skin		(2 marks)




Rr



- Genotypic ratio (i) 1 2 1 (1 mark)
- Phenotypic ratio (ii) 3 smooth coats 1 wrinkled coat; (1 mark)
- (c) The total number of wrinkled seeds.

$$1/4 \times 14,640 = 3660$$
; (2 marks)

5.	(a)	(i)	 It is long/wide/broad/flat; to provide a large surface area for attachment of muscles; 	
			- Has facets; for articulation with sacrum;	(2 marks)
		(ii)	J Has flexible cartilage; which allows for widening of the (female) pel girdle when giving birth/to absorb shock.	vic
				(2 marks)
	(b)	Allows	s passage of blood vessels/nerves/ and muscles;	(1 mark)
	(c)	(i) (ii)	Femur; Ball and socket;	1 mark 1 mark
	(d)	Coccy	x;	1 mark
6.	(a)	See gra	aph on page 5.	
	(b)	(i)	No change in population/population is constant; mice still maturing/have not given birth;	
				(2 marks)
		(ii)	Slow/gradual population growth; few mice have reached sexual maturity;	(2 marks)
		(iii)	Faster/rapid rate of population growth/exponential; Many mice sexually matured/reproducing/enough food/space/no competition/ birth rate higher than death/no diseases:	(2 marks)
		(iv)	Population decline; Competition is high / food is limiting / space is limiting/accumulation of toxic waste/disease (outbreak) deathrate higher than birth rate.	
				(2 marks)
	(c)	(i)	6 and 8;	(1 mark)
		(ii)	310 - 115 = 195 mice per month;	(2 marks)
	(d)	Popula	ation would increase;	(1 marks)
	(e)	Food;	space ; cage size; water;	
			(max)) (2 marks)



(a) When a blood vessel is cut/injured platelets/thrombocytes/damaged tissue/wound is exposed to the air; they release thrombokinase/thromboplastin; an enzyme that activates the conversion of prothrombin; to thrombin; in the presence of calcium ions; vitamin K/ phylloquinone; is needed for the formation of prothrombin; Thrombin converts (soluble blood protein) fibrinogen; into (the fibrous form) fibrin; which forms a mesh / network across the wound; The clot so formed prevents excessive bleeding; and entry of disease agents/pathogens/micro-organisms/microbes;

7.

Max 10 marks

- (b) Many to provide a large surface area; across which large amounts of gases diffuse; moist surfaces; to dissolve respiratory gases; so as to diffuse. Made of a thin membrane/epithelium/one cell thick wall; to reduce diffusion distance; Highly vascularized; to carry away oxygen; and bring in carbon (IV) oxide; creating a steep diffusion gradients. (10 marks)
- 8. (a) Regulation of blood sugar; when blood sugar is below normal/90 mg/100 cm³ glucagon; triggers the conversion of glycogen to glucose in the liver; the glucose is released into the blood stream. When blood sugar is in excess above normal/10 mg/100 cm³, insulin; causes the liver to convert glucose excess to glycogen; which is stored.

Production of heat energy; by increasing the rate of metabolic activities;

Excretion of bile pigments; produced due to breakdown of worn out red blood cells; Deamination/removal of amino group of excess amino acids to form urea; and detoxication/poisonous/toxic substances;

(Max 10 marks)

(b) Sweat glands excrete urea; excess water; and salts; hence maintaining salt & water balance in the blood. Evaporation of sweat; cools the body due to loss of latent heat of vaporization; when the body temperature rises; blood vessels in the skin vasolidate; allowing more blood to flow near the skin surface; thus heat is lost to the environment by radiation/convection. The erctor pili mucle relaxes hair flattens; in a hot environment reducing insulation; hence heat is lost from the body by radiation/convection; to the environment.

(max 10 marks)

4.4.3 Biology Paper 3 (231/3)

- **1.** (a) (i) Sternum; (1 mark)
 - (ii) The internal intercostal muscles relax; pulling the ribs upwards; and outwards; This increases the volume of the rib cage while pressure decreases; Forcing air into the lungs;

(5 marks)

- (b) (i) Anterior/dorsal view; (1 mark)
 - (ii) Name Neural canal; (1 mark)

Function - Passage of the spinal cord. (1 mark)

(iii) V: It is thick and solid; for bearing the weight of the body (back) (2 marks)

 ${f S}\,:\,$ It is long; to provide a large surface area for attachment of muscles;

(2 marks)

- (c) (i) Image width = 9.8 cm;
 - (ii) Magnification = $\frac{\text{Image length / width}}{\text{Actual length / width}}$;

$$=$$
 $\frac{9.8 \pm 0.1}{4.6 \pm 0.1}$

 $Mg = \times 2.13$;

(iii) Actual length AB = $\frac{10.4 \pm 0.1}{2.13}$;

= 4.8826 cm;

(5 marks)

Food Substance Tested	Procedure	Observation	Conclusion
Reducing sugars	 Put 2 cm³ of C in a test tube; Add equal volume of Benedict's blu Solution. Put in a hot water bath/heat/ warm/boil; 	No colour change/ e colour remains/ colour of Benedict's solution remains/ persists;	Reducing sugars absent;
2. Reducing sugar	 Put 2 cm ³ of C in a test tube; Add a few drops of dilute hydrochloric acid. Place the test tube in a hot water bath for 3 minutes; Remove the test tube and cool in cold water. Add (NaH) 2CO 3 drop by drop until fizzing stops Add 2 cm 3 of Benedict's Solution. Place the test tube in a hot water bath/heat/warm/boil; 	Colour changes to green / yellow / orange / brown;	Reducing sugars present;
3. Proteins	 Put 2 cm ³ of C in a test tube; Add an equal amount of sodium hydroxide solution and shake. Add copper sulphate drop by drop, shaking well after each addition; 	Colour changes to purple/violet/mauve;	Proteins present;

3.

1.	(a)	Simple leaves	go to 2;
	(b)	Compound leaves	go to 4;
2.	(a)	Leaves net-veined/reticulate	go to 3;
	(b)	Leaves parallel veined	Commelinaceae;
3.	(a)	Leaves with serrated margins	Malvaceae;
	(b)	Leaves with smooth (entire) margins	Nystaginaceae;
4.	(a)	Leaves opposite	go to 5;
	(b)	Leaves alternate	Bignoniceae;
5.	(a)	Leaves pinnate	Papilionaceae;
	(b)	Leaves trifoliate	Compositae;
			(10 marks)