

NAME:INDEX NO:.....

SCHOOL:SIGNATURE :.....

DATE:

233/3

Chemistry

(Practical)

July/August 2016

Time: 2 Hours

KAKAMEGA SOUTH SUB-COUNTY JOINT EVALUATION TEST – 2016

Kenya Certificate of Secondary Examination (KCSE)

233/3

Chemistry

(Practical)

INSTRUCTIONS TO CANDIDATES

1. Write your name and index number in the spaces provided above.
2. Sign and write the date of examination in the spaces provided above.
3. You are not allowed to work with apparatus for the first 15 minutes of the 2 ¼ hours for this paper. This time is to enable you read the question paper and make sure you have all chemicals and apparatus that you need.
4. All workings MUST be clearly shown.
5. KNEC Mathematical tables and silent electronic calculators may be used.

QUESTION	MAXIMUM SCORE	CANDIDATE'S SCORE
1	19	
2	7	
3	14	
TOTAL	40	

This paper consists of 8 printed pages Check the Question paper to ensure that all pages are printed as indicated and no question are missing.

1. You are provided with:
- Solution P. 0.2m Hydrochloric acid.
 - Solution Q sodium hydroxide solution
 - Solution R, Containing 49g/l of dibasic acid H₂A

You are required to:-

- Dilute Solution Q with distilled water.
- Standardize the diluted solution Q with solution P
- Determine the relative formula mass of A.

Procedure I

- Pipette 25 cm³ of solution Q into a clean dry 250cm³ volumetric flask.
- Measure 175 cm³ of distilled water using a 100cm³ measuring cylinder and add to solution Q in the flask.
- Shake well.
- Label this as solution S and keep it for further tests in procedure II.
- Pipette 25cm³g solution S into a clean dry conical flask.
- Add 2 to 3 drops of phenolphaclein indicator and titrate with solution P.
- Record your results in the table I below.
- Repeat the procedure to obtain accurate results.

Table I

(4mks)

	1	2	3
Final Burette reading (cm ³)			
Initial burette reading (cm ³)			
Volume of solution P used (cm ³)			

1. a) Determine the average volume of solution P used. (1mk)

.....

- b) i) Find the number of moles of solution P used to react with 25cm³ of the diluted solution S. (1mk)

.....

- ii) Find the numbers of moles of solution S in 25 cm^3 of diluted solution. (1mk)
- iii) Determine the number of moles of sodium hydroxide contained in the 1000 cm^3 of solution S. (1mk)
- c) Using your results in b (ii) above determine the concentration in moles per litre of the original sodium hydroxide solution Q (1mk)

Procedure II

- Pipette 25 cm^3 of the standardized solution into a clean dry conical flask.
- Empty your burette completely off solution P.

Now fill your burette with solution R and titrate with solution S in the conical flask containing 2 to 3 drops of phenolphthalein indicator.

- Record your results in table II below.
- Repeat the procedure to obtain accurate results.

TABLE II	1	2	3
Final burette reading (cm ³)			
Initial burette reading (cm ³)			
Volume of solution R used (cm ³)			

(4mks)

- d) Determine the average volume of solution R used. (1mk)

.....

- e) Determine the number of moles of sodium hydroxide in 25 cm³ of solution S and hence the number of moles of solution R used. (2mks)

.....

- f) Find the number of moles of solution R contained in one litre of solution. (1mk)

.....

- g) Given that relative atomic mass of H= 1.0,

- i) Find the relative formula mass of re dibasic acid. H₂A. (1mk)

- ii) Determine the relative formula mass of A in the formula H₂A (1mk)

Q2. FA₁ is 3.5g of a metal carbonate XCO₃ FA₂ is 2M HCl

The carbonate and acid react according to the following equation.



The enthalpy change, ΔH for this reaction is $-59.5 \text{ KJ mol}^{-1}$. You're required to determine to determine the temperature rise when a known mass of one solid carbonate. CXO₃ is added to an excess of hydrochloric acid and use your results to calculate atomic mass R.A.M of the metal X.

Procedure.

Using the measuring cylinder provided add 50 cm³ of hydrochloric acid FA₂ to 250 cm³ plastic beaker.

- Measure the initial temperature of the acid FA₂ In the plastic beaker and record it in the table below.
- Empty all the carbonate provided FA₁ into the acid and stir gently with the thermometer.
- Record the maximum temperature attained when the solid has reacted with the acid in the table below.

Maximum temperature attained (°C)	
Initial temperature of FA ₂ (°C)	
Temperature change ΔH °C	

i) Using the temperature change from (a) above.

Calculate the enthalpy change for the reaction.

(Specific heat capacity of a solution = 4.2J/g/K, density of solution = 1g/cm³ (2mks)

ii) Using your answer in (i) above and ΔH value for the reaction. Calculate the number of moles of FA₁ that reacted. (1mk)

- iii) Using the mass of FA₁ given calculate the relative atomic mass RAM of metal X
 (C= 12.0, O= 16.0) (1mk)

Q3. You are provided with solid H. Carry out the tests below and write your observations and inferences in the spaces provided.

- a) Place about half of solid H in a clean dry test tube. Heat the solid gently and then strongly. Test for any gas produced using both the red and blue litmus paper.

Observation	Inferences
(1mk)	(1mk)

- b) Dissolve the remaining portion g solid H in about 8 cm³ of distilled water contained in a boiling tube. Divide the solution into three portions.

- i) To the first portion add aqueous solution hydroxide drop wise until in excess.

Observation	Inferences
	(½ mk)

1 mk

- ii To the second portion add two drops of concentrated nitric (v) acid then followed by aqueous sodium hydroxide dropwise until in excess.

Observation	Inferences
(1mk)	(1mk)

- iii) To the third portion add 2-3 drops of barium chloride solution.

Observation	Inferences
(1mk)	(1mk)

- iv) To the mixture obtained in (b iii) above add 2 cm³ aqueous hydrochloride acid.

Observation	Inferences
(¹ / ₂ mk)	(¹ / ₂ mk)

- C) You are provided with substance v. Carry out the following test and record your observation and inferences in the spaces provided.

- i) Place about a half of substance V on metallic spatula and ignite it a Bunsen burner flame.

Observation	Inferences
(1mk)	(1mk)

- ii) Place the remaining amount of substance V in a clean boiling tube add about 10 cm³ of distilled water. Warm the mixture for about 10 seconds and divide it into two portions.

- i) To the first portion add 2 to 3 drops of universal indicator.

Observation	Inferences
(1mk)	(1mk)

- ii) To the second portion add 2 to 3 drops of acidic potassium manganate (VII).

Observation	Inferences
(1mk)	(1mk)