




		Working Space
3.	<p><b>1990 Q8 P2</b></p> <p>In a triangle ABC, D is the midpoint of AB and E is a point on BC such that <math>BE = \frac{2}{3} BC</math>. If <math>\mathbf{AD} = \mathbf{p}</math> and <math>\mathbf{AC} = \mathbf{q}</math>, express <math>\mathbf{EC}</math> in terms of <math>\mathbf{p}</math> and <math>\mathbf{q}</math>.</p> <p style="text-align: right;">(2 marks)</p>	
4.	<p><b>1990 Q10 P2</b></p> <p>A point T divides a line AB internally in the ratio 5 : 2. Given that A is (-4, 10) And B is (10, 3) find the coordinates of T.</p> <p style="text-align: right;">(4 marks)</p>	
5.	<p><b>1991 Q6 P1</b></p> <p>In the diagram below OABC is a parallelogram.</p>  <p>AB is produced to T such that <math>BT:AB = 1:2</math>. M is the midpoint of AC. Given that <math>\mathbf{OA} = \mathbf{a}</math> and <math>\mathbf{OC} = \mathbf{c}</math>. Express</p>	




		Working Space
8.	<p><b>1992 Q24 P1</b></p> <p>OABC is a trapezium such that the coordinates of O,A,B and C are (0,0),(2,-1), (4, 3) and (0, y).</p> <p>a) Find the value of y (2 marks)</p> <p>M is a midpoint of AB and N is a midpoint of OM. Show that A, N and C are collinear. (6 marks)</p>	
9.	<p><b>1992 Q7 P2</b></p> <p>The vectors <b>p</b>, <b>q</b> and <b>y</b> are expressed in terms of the vectors <b>t</b> and <b>s</b> as follow:</p> $\mathbf{p} = 3\mathbf{t} + 2\mathbf{s}$ $\mathbf{q} = 5\mathbf{t} - \mathbf{s}$ $\mathbf{y} = h\mathbf{t} + (h - k)\mathbf{s}$ <p>where <b>h</b> and <b>k</b> are constants. Given that <math>\mathbf{y} = 2\mathbf{p} - 3\mathbf{q}</math>, find the values of <b>h</b> and <b>k</b>. (4marks)</p>	
10	<p><b>1993 Q21 P1</b></p> <p>OABC is a trapezium in which <b>OA = a</b>, <b>OC = c</b> and <b>CB = 3a</b>. CB is produced to such that <b>CB : BD = 3: 1</b>. E is a point on AB such that <b>AB = 2AE</b>. Show that O, E and d are collinear. (8 marks)</p>	

working space

11 **1993 Q16 P1**


In the figure below  $CA = b$   $CB = a$ ,  $AX = XY$  and  $AY = YB$ .



**A**      **X**      **Y**      **B**  
Express  $CX$  in terms of  $a$  and  $b$       (3 marks)

12 **1994 Q24 P1**

In the figure below  $AB = a$ ,  $AD = b$ ,  $AX : XC = 2:3$  and  $XB = 4:5$




	<p style="text-align: center;">A                  b                  D</p> <p>a) Express</p> <p>  i) <b>AC</b></p> <p>  ii) <b>DC</b> in terms of <b>a</b> and <b>b</b> in the simplest form. (6 marks)</p> <p>b) If <b>DC = na + mb</b>, find the values of <i>n</i> and <i>m</i> (2 marks)</p>	Working Space
13	<p><b>1994Q12P2</b></p> <p>Find the position vector of point R which divides line MN internally in the ratio 2: 3. Take the position vectors of M and N to be</p> $\mathbf{M} = \begin{pmatrix} 3 \\ 4 \\ -6 \end{pmatrix} \quad \text{and } \mathbf{N} = \begin{pmatrix} 3 \\ 4 \\ -6 \end{pmatrix}$ $\begin{pmatrix} -5 \\ -1 \\ 2 \end{pmatrix}$ <p style="text-align: right;">(3 marks)</p>	
14	<p><b>1994 Q10 P2</b></p> <p>In the figure below <math>OC = 3 CA</math> and <math>OD = 3DB</math>. By taking <math>OA = a</math>, <math>OB = b</math>, show that <math>CD \parallel AB</math>. (3 marks)</p> <p style="text-align: center;">O</p>	

	<p>D                      C                                      D</p> <p>B                      A                                      B</p>	
--	---	--

15	<p><b>1994 Q15 P2</b>          In the figure below ABCD is a parallelogram. AOC and BOD are diagonals of the parallelogram. Show that the diagonals of the parallelogram bisect each other. Give reasons. (3 marks)</p> <p style="text-align: center;">A                                      B</p> <p style="text-align: center;">D                                      C</p>	Working Space
----	---	---------------


16	<p><b>1995 Q 18 P1</b>          The figure below is a right pyramid with a rectangular base ABCD and VO as the height. The vectors <math>\mathbf{AD} = \mathbf{a}</math>, <math>\mathbf{AB} = \mathbf{b}</math> and <math>\mathbf{DV} = \mathbf{c}</math></p> <p>a) Express (i) <math>\mathbf{AV}</math> in terms of <math>\mathbf{a}</math> and <math>\mathbf{c}</math> (1 mark)</p>	
----	---	--

	<p>(ii) <math>\mathbf{BV}</math> in terms of <math>\mathbf{a}</math>, <math>\mathbf{b}</math> and <math>\mathbf{c}</math> ( 2 marks)</p> <p>(b) M is point on <math>\mathbf{OV}</math> such that <math>\mathbf{OM} : \mathbf{MV} = 3:4</math>, Express <math>\mathbf{BM}</math> in terms of <math>\mathbf{a}</math>, <math>\mathbf{b}</math> and <math>\mathbf{c}</math>. Simplify your answer as far as possible ( 5 marks)</p>	
<p>17</p>	<p><b>1996 Q 22 P1</b></p> <p>a) In the diagram below <math>OABC</math> is a parallelogram, <math>\mathbf{OA} = \mathbf{a}</math> and <math>\mathbf{AB} = \mathbf{b}</math>. <math>N</math> is a point on <math>\mathbf{OA}</math> such that <math>\mathbf{ON} : \mathbf{NA} = 1 : 2</math></p>  <p>(b) Find</p> <p>(i) <math>\mathbf{AC}</math> in terms of <math>\mathbf{a}</math> and <math>\mathbf{b}</math></p> <p>(ii) <math>\mathbf{BN}</math> in terms of <math>\mathbf{a}</math> and <math>\mathbf{b}</math></p> <p>(c) The lines <math>AC</math> and <math>BN</math> intersect at <math>X</math>, <math>\mathbf{AX} = h\mathbf{AC}</math> and <math>\mathbf{BX} = k\mathbf{BN}</math></p> <p>(i) By expressing <math>\mathbf{OX}</math> in two ways, find the values of <math>h</math> and <math>k</math></p> <p>(ii) Express <math>\mathbf{OX}</math> in terms of <math>\mathbf{a}</math> and <math>\mathbf{b}</math> (1 mark)</p>	<p>Working Space</p>
<p>18</p>	<p><b>1997 Q 11 P2</b></p> <p><math>ABC</math> is a triangle and <math>P</math> is on <math>AB</math> such that <math>P</math> divides <math>AB</math> internally in the ratio <math>4:3</math>. <math>Q</math> is a point on <math>AC</math> such that <math>PQ</math> is parallel to <math>BC</math>. If <math>AC = 14</math> cm</p> <p>(i) State the ratio <math>AQ:QC</math></p> <p>(ii) Calculate the length of <math>QC</math></p>	



19 1997 Q 22 P1

In the figure below  $OA = a$ ,  $OB = b$ ,  $AB = BC$  and  $OB:BD = 3:1$




- (a) Determine
- (i)  $AB$
  - (ii)  $CD$ , in terms of  $a$  and  $b$
- (b) If  $CD : DE = 1:k$  and  $OA:AE = 1: m$  determine
- (i)  $DE$  in terms of  $a$ ,  $b$  and  $k$

Working Space

20 **1998 Q 9 P2**

In the figure, KLMN is a trapezium in which KL is parallel to NM and  $KL = 3 NM$




Given that  $KN = w$ ,  $NM = u$  and  $ML = v$ . Show that  $2u = v + w$

21 **1998 Q 22 P1**

The figure below shows a grid of equally spaced parallel lines

$AB = a$  and  $BC = b$




Working Space

	<p>(a) Express</p> <p>(i) <b>AC</b> in terms of a and b</p> <p>(ii) <b>AD</b> in terms of a and b.</p> <p>(b) Using triangle BEP, express <b>BP</b> in terms of a and b</p> <p>(c) PR produced meets BA produced at X and  <math>\mathbf{PR} = \frac{1}{9}\mathbf{b} - \frac{8}{3}\mathbf{a}</math></p> <p>By writing <b>PX</b> as <math>k\mathbf{PR}</math> and <b>BX</b> as <math>h\mathbf{BA}</math> and using the triangle BPX determine the ratio PR: RX</p>	
22	<p><b>1999 Q 14 P2</b></p> <p>The points P, Q and R lie on a straight line. The position vectors of P and R are <math>2\mathbf{i} + 2\mathbf{j} + 13\mathbf{k}</math> and <math>5\mathbf{i} - 3\mathbf{j} + 4\mathbf{k}</math> respectively. Q divides PR Internally in the ratio 2:1. Find the</p> <p>(a) Position vector of Q.</p> <p>(b) Distance of Q from the origin</p>	
23	<p><b>1999 Q 21 P1</b></p> <p>In triangle OAB, <math>\mathbf{OA} = \mathbf{a}</math>, <math>\mathbf{OB} = \mathbf{b}</math> and P lies on AB such that AP: BP = 3:5</p> <p>(a) Find the terms of <b>a</b> and <b>b</b> the vectors</p> <p>(i) <b>AB</b></p> <p>(ii) <b>AP</b></p> <p>(iii) <b>BP</b></p> <p>(iv) <b>OP</b></p> <p>(b) Point Q is on OP such <math>AQ = \frac{-5}{8}\mathbf{a} + \frac{9}{40}\mathbf{b}</math>.</p> <p>Find the ratio OQ: QP</p>	

24 **2000 Q 21 P1**

The figure below shows triangle OAB in which M divides OA in the ratio 2: 3 and N divides OB in the ratio 4:1 AN and BM intersect at X.



(a) Given that  $OA = \mathbf{a}$  and  $OB = \mathbf{b}$ , express in terms of  $\mathbf{a}$  and  $\mathbf{b}$ :

(i)  $\mathbf{AN}$

(ii)  $\mathbf{BM}$

(b) If  $\mathbf{AX} = s \mathbf{AN}$  and  $\mathbf{BX} = t \mathbf{BM}$ , where  $s$  and  $t$  are constants, write two expressions for  $\mathbf{OX}$  in terms of  $\mathbf{a}, \mathbf{b}$ ,  $s$  and  $t$ . Find the value of  $s$ . Hence write  $\mathbf{OX}$  in terms of  $\mathbf{a}$  and  $\mathbf{b}$ .


25 **2001 Q 16 P1**

The position vectors for points P and Q are  $4\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$  and  $3\mathbf{i} - 6\mathbf{j} + 6\mathbf{k}$  respectively. Express vector  $\mathbf{PQ}$  in terms of unit vectors  $\mathbf{i}$ ,  $\mathbf{j}$  and  $\mathbf{k}$ . Hence find the length of  $\mathbf{PQ}$ , leaving your answer in simplified form.

26

**2001 Q 19 P1**

The figure below shows a parallelogram OPQR with O as the origin,  $\mathbf{OP} = \mathbf{p}$  and  $\mathbf{OR} = \mathbf{r}$ , Point T divides RQ in the ratio 1:4 and PT meets OQ at S.



- (a) Express in terms of  $\mathbf{p}$  and  $\mathbf{r}$  the vectors
- $\mathbf{OQ}$
  - $\mathbf{OT}$
- (b) Vector  $\mathbf{OS}$  can be expressed in two ways:  $m\mathbf{OQ}$  or  $\mathbf{OT} + n\mathbf{TP}$ , Where  $m$  and  $n$  are constants express  $\mathbf{OS}$  in terms of
- $m$ ,  $\mathbf{p}$  and  $\mathbf{r}$
  - $n$ ,  $\mathbf{p}$  and  $\mathbf{r}$
- Hence find the:
- Value on  $m$
  - Ratio OS: SQ

27


**2002 Q 10 P2**

The coordinates of points O,P,Q and R are  $(0,0)$ ,  $(3,4)$ ,  $(11,6)$  and  $(8,2)$  respectively. A point T is such that vectors  $\mathbf{OT}$ ,  $\mathbf{QP}$  and  $\mathbf{QR}$  satisfy the vector equation.  $\mathbf{OT} = \mathbf{QP} + \frac{1}{2}\mathbf{QR}$ . Find the coordinates of T.

28

**2002 Q 4 P1**

The position vectors of points X and Y are  $\mathbf{x} = 2\mathbf{i} + \mathbf{j} - 3\mathbf{k}$  and  $\mathbf{y} = 3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$  respectively. Find  $\mathbf{XY}$

		Working Space
29	<p><b>2003 Q 6 P1</b>  Given that <math>x = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}</math>, <math>y = -3\mathbf{i} + 4\mathbf{j} - \mathbf{k}</math> and <math>z = -5\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}</math> and that <math>\mathbf{p} = 3\mathbf{x} - \mathbf{y} + 2\mathbf{z}</math>. Find the magnitude of vector <math>\mathbf{p}</math> to 3 significant figure (4marks)</p>	
30	<p><b>2003 Q 21 P1</b>  In the figure below, vector <math>OP = \mathbf{p}</math> and <math>OR = \mathbf{r}</math>. Vector <math>OS = 2\mathbf{r}</math> and <math>OQ = \frac{3}{2}\mathbf{p}</math>.</p>  <p>a) Express in terms of <math>\mathbf{p}</math> and <math>\mathbf{r}</math> (i) <math>\mathbf{QR}</math> and (ii) <math>\mathbf{PS}</math>  b) The lines <math>QR</math> and <math>PS</math> intersect at <math>K</math> such that <math>\mathbf{QK} = m \mathbf{QR}</math> and <math>\mathbf{PK} = n \mathbf{PS}</math>, where <math>m</math> and <math>n</math> are scalars. Find two distinct expressions for <math>\mathbf{OK}</math> in terms of <math>\mathbf{p}</math>, <math>\mathbf{r}</math>, <math>m</math> and <math>n</math>. Hence find the values of <math>m</math> and <math>n</math>. (5marks)  c) State the ratio <math>PK:KS</math></p>	
31	<p><b>2004 Q 4 P1</b>  Given that <math>\mathbf{OA} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}</math> and <math>\mathbf{OB} = 4\mathbf{i} + \mathbf{j} - 3\mathbf{k}</math>. Find the distance between points <math>A</math> and <math>B</math> to 2 decimal places.</p>	

		Working Space
32	<p><b>2004 Q 21 P1</b></p> <p>a) If A, B and C are the points P and Q are p and q respectively is another point with position vector <math>r = \frac{3}{2}q - \frac{1}{2}p</math>. Express in terms of p and q.</p> <p>i) <b>PR</b></p> <p>ii) <b>RQ</b> hence show that P, Q and R are collinear.</p> <p>iii) Determine the ratio PQ: QR.</p>	
33	<p><b>2005 Q 13 P1</b></p> <p>Point T is the midpoint of a straight line AB. Given the position vectors of A and T are <math>i-j+k</math> and <math>2i+1\frac{1}{2}k</math> respectively, find the position vector of B in terms of i, j and k. ( 3 marks)</p>	
34	<p><b>2005 Q 18 P1</b></p> <p>The points P, Q, R and S have position vectors <math>2p, 3p, r</math> and <math>3r</math> respectively, relative to an origin O. A point T divides PS internally in the ratio 1:6</p> <p>(a) Find, in the simplest form, the vectors <b>OT</b> and <b>QT</b> in terms <b>P</b> and <b>r</b> ( 4 marks)</p> <p>(b) (i) Show that the points Q, T, and R lie on a straight line ( 3 marks)</p> <p>(ii) Determine the ratio in which T divides QR ( 1 mark)</p>	

working space

35 **2006 Q 12 P1**

Two points P and Q have coordinates (-2, 3) and (1,3) respectively. A translation map point P to P' ( 10, 10)

a) Find the coordinates of Q' the image of Q under the translation ( 1 mark)

(ii) The position vector of P and Q in (a) above are p and q respectively given that  $m\mathbf{p} - n\mathbf{q} =$


$$\begin{pmatrix} -12 \\ 9 \end{pmatrix}$$

( 3 marks)

b) Find the value of m and n

36 **2006 Q 22 P1**


In the diagram below, the coordinates of points A and B are (1,6) and (15,6) respectively). Point N is on OB such that  $3 ON = 2OB$ . Line OA is produced to L such that  $OL = 3 OA$




(a) Find vector LN

( 3 marks)



	<p>(b) Given that a point M is on LN such that <math>LM:MN = 3:4</math>, find the coordinates of (2 marks)</p> <p>(c) If line OM is produced to T such that <math>OM:MT = 6:1</math></p> <p>(i) Find the position vector of T (1 mark)</p> <p>(ii) Show that points L, T and B are collinear (4 marks)</p>	Working Space
37	<p><b>2006 Q 9 P2</b></p> <p>Given that <math>q \mathbf{i} + \frac{1}{3} \mathbf{j} + \frac{2}{3} \mathbf{k}</math> is a unit vector, find q (2 marks)</p>	
38	<p><b>2007 Q 21 P1</b></p> <p>In the figure below, <math>\mathbf{OQ} = q</math> and <math>\mathbf{OR} = r</math>. Point X divides OQ in the ratio 1: 2 and Y divides OR in the ratio 3: 4 lines XR and YQ intersect at E.</p>  <p>(a) Express in terms of q and r</p> <p>(i) <math>\mathbf{XR}</math> (1 mark)</p> <p>(ii) <math>\mathbf{YQ}</math> (1 mark)</p> <p>(b) If <math>\mathbf{XE} = m \mathbf{XR}</math> and <math>\mathbf{YE} = n \mathbf{YQ}</math>, express <math>\mathbf{OE}</math> in terms of: (1 mark)</p> <p>(i) r, q and m</p> <p>(ii) r, q and n (1 mark)</p> <p>(c) Using the results in (b) above, find the values of m and n. (6 marks)</p>	

39	<p><b>2007 Q 12 P2</b>          Vector <math>q</math> has a magnitude of 7 and is parallel to vector <math>p</math>.          Given that <math>p = 3\mathbf{i} - \mathbf{j} + 1\frac{1}{2}\mathbf{k}</math>, express vector <math>q</math> in terms of <math>\mathbf{i}</math>, <math>\mathbf{j}</math>, and <math>\mathbf{k}</math>.          ( 2 marks)</p>	Working Space
40	<p><b>2008 Q 19 P2</b>          In the figure below <math>AB = \mathbf{p}</math>, <math>AD = \mathbf{q}</math>, <math>DE = \frac{1}{2} AB</math> and <math>BC = \frac{2}{3} BD</math></p>  <p>a) Find in terms of <math>\mathbf{p}</math> and <math>\mathbf{q}</math> the vectors: (1mark)</p> <p>(i) <math>\mathbf{BD}</math>;          (1mark)</p> <p>(ii) <math>\mathbf{BC}</math>;          (1mark)</p> <p>(iii) <math>\mathbf{CD}</math>;          (1mark)</p> <p>(iv) <math>\mathbf{AC}</math>. (2marks)</p> <p>b) Given that <math>\mathbf{AC} = k\mathbf{CE}</math>, where <math>k</math> is a scalar, find</p> <p>(i) The value of <math>k</math> (4marks)</p> <p>(ii) The ratio in which C divides AE (1mark)</p>	

41	<p><b>2008 Q 4 P2</b></p> <p>The position vectors of points A and B are <math>\begin{pmatrix} 3 \\ -1 \\ -4 \end{pmatrix}</math> and <math>\begin{pmatrix} 8 \\ -6 \\ 6 \end{pmatrix}</math> respectively.</p> <p>A point P divides AB in the ratio 2:3. Find the position Vector of point P. (3marks)</p>	
42	<p><b>2009 Q 20 P1</b></p> <p>The position vectors of point A and B with respect to the O, are <math>\begin{pmatrix} -8 \\ 5 \end{pmatrix}</math> and <math>\begin{pmatrix} 12 \\ -5 \end{pmatrix}</math> respectively</p> <p>Point M is the midpoint of AB and N is the midpoint of OA.</p>	Working Space
	<p>(a) Find:</p> <p>i) The coordinates of N and M ( 3 marks)</p> <p>ii) The magnitude of NM ( 3 marks)</p> <p>(b) Express vector <b>NM</b> in term of <b>OB</b>.</p> <p>(c) Point P maps onto P' by a translation <math>\begin{pmatrix} -5 \\ 8 \end{pmatrix}</math></p> <p>Given that <b>OP=OM+2MN</b>, find the coordinates of P'</p>	

43 **2009 Q 6 P2**


Vector  $\mathbf{OA} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$  and  $\mathbf{OB} = \begin{pmatrix} 6 \\ -3 \end{pmatrix}$  Point C is on

OB such  $CB=2OC$  and point D is on AB such that  $AD=3DB$ .

Express  $\mathbf{CD}$  as a column vector. (3 marks)


44 **2010 Q 7 P1**

In the figure below, OPQR is a trapezium in which PQ is parallel to OR and M is the mid-point of QR and  $\mathbf{OP} = \mathbf{p}$ ,  $\mathbf{OR} = \mathbf{r}$  and  $\mathbf{PQ} = \frac{1}{3}\mathbf{OR}$ .



Find  $\mathbf{OM}$  in terms of  $\mathbf{p}$  and  $\mathbf{r}$ . (3 marks)

Working Space

<p>45</p>	<p><b>2010 Q 18 P2</b>          In the figure below OJKL is a parallelogram in which  <math>OJ = 3p</math> and <math>OL = 2r</math></p>  <p>a) If A is a point on LK such that <math>LA = \frac{1}{2} AK</math> and point B divide the line JK externally in the ratio 3:1, express <b>OB</b> and <b>AJ</b> in terms of <b>p</b> and <b>r</b>.          (2 marks)</p> <p>b) Line OB interests AJ at X such that <b>OX</b> = <b>mOB</b> and <b>AX</b> = <b>nAJ</b>.</p> <p>i) Express <b>OX</b> in terms of <b>p</b>, <b>r</b> and <b>m</b>. (1 mark)          ii) Express <b>OX</b> in terms of <b>p</b>, <b>r</b> and <b>n</b> (1 mark)          iii) Determine the value of <b>m</b> and <b>n</b> and hence the ratio in which point x divides line AJ.          (6 marks)</p>	
<p>46</p>	<p><b>2011 Q 13 P2</b>          Vector <b>OP</b>=<math>6\mathbf{i} + \mathbf{j}</math> and <b>OQ</b>+ <math>-2\mathbf{i} + 5\mathbf{j}</math>. A point N divides <b>PQ</b> internally in the ratio 3:1. Find <b>PN</b> in terms of <b>i</b> and <b>j</b>.          ( 3 marks)</p>	



